

Medium effects in strange quark matter and strange stars*

K. Schertler, C. Greiner, and M.H. Thoma[†]
Institut für Theoretische Physik, Universität Giessen,
35392 Giessen, Germany

November, 1996

Abstract

We investigate the properties of strange quark matter at zero temperature including medium effects. The quarks are considered as quasiparticles which acquire an effective mass generated by the interaction with the other quarks of the dense system. The effective quark masses are derived from the zero momentum limit of the dispersion relations following from an effective quark propagator obtained from resumming one-loop self energy diagrams in the hard dense loop approximation. This leads to a thermodynamic selfconsistent description of strange quark matter as an ideal Fermi gas of quasiparticles. Within this approach we find that medium effects reduce the overall binding energy with respect to ^{56}Fe of strange quark matter. For realistic values of the strong coupling constant strange quark matter is not absolutely stable. The application to pure strange quark matter stars shows that medium effects have, nevertheless, no impact on the mass-radius relation of the stars. However, a phase transition to hadronic matter at the surface of the stars becomes more likely.

PACS: 12.38.Mh, 26.60.+c

Keywords: Strange quark matter, effective masses, quark stars

1 Introduction

Strange quark matter (SQM) or strangelets are thought to be confined (bulk) objects containing a large number of delocalized quarks ($u\dots u$, $d\dots d$, $s\dots s$), so-called multi-quark droplets. Multi-quark states consisting only of u - and d -quarks must have a mass larger than ordinary nuclei, otherwise normal nuclei would certainly be unstable, which, of course, is not the case. However, the situation is different for SQM, which would contain approximately the same amount of u -, d - and s -quarks. It has been suggested as a possible absolutely stable or metastable phase of nuclear matter [1], which might be realized in form of small droplets (strangelets) or in the interior of superdense stars [2]. The proposal that hypothetical SQM at zero temperature and in β -equilibrium might be absolutely stable has stimulated substantial activity. Such a scenario would be of fundamental importance, as, for example, it could explain a large fraction of the non-observable mass of the universe in certain cosmological models, and could modify substantially our understanding of the structure and stability of neutron stars [2]. The reason for the possible stability lies in introducing a third flavor degree of free-

dom, the strangeness, where the mass of the strange quarks is considerably smaller than the Fermi energy of the quarks, thus lowering the total mass per unit baryon number of the system. The equation of state of SQM has been described as a non-interacting gas of quarks at zero temperature, taking into account the phenomenological bag constant. Also quark interactions within lowest order perturbative QCD have been considered [3, 4]. First studies in the context of the MIT-bag model predicted that sufficiently heavy strangelets might be absolutely stable [5], i.e., that the energy per baryon might be lower than the one of ^{56}Fe . On the other hand, for smaller strangelets it is also conceivable that the mass per baryon is lower than the mass of the strange Λ -baryon, but larger than the nucleon mass. The droplet is then in a metastable state, i.e., it cannot decay (strongly) into nonstrange hadronic matter [4]. Metastable strangelets relax somewhat the stringent conditions on the choice of parameters of the bag model required by an absolutely stable state [5], namely small bag constants $B_0^{1/4} \lesssim 150$ MeV. According to this picture, SQM should appear as a nearly neutral

*Work supported by BMBF, GSI Darmstadt, and DFG

[†]Heisenberg fellow

and massive state because the number of strange quarks is nearly equal to the number of massless up or down quarks and so the strange quarks neutralize that hypothetical form of nuclear matter.

In condensed matter as well as nuclear physics medium effects play an important role. One of the most important medium effects are effective masses generated by the interaction of the particles with the system. In a gluon gas at high temperature the consideration of an effective mass for the gluons within the ideal gas approximation lead to an excellent description of the equation of state found in lattice calculations [6, 7]. The effective gluon mass has been derived from the gluon self energy, containing the interaction of the gluons with the heat bath. Using the high temperature approximation [8] or, equivalently, the hard thermal loop (HTL) approximation [9], which ensures a gauge invariant treatment, a gluon self energy of the order $g^2 T^2$ has been found. Resumming this self energy leads to an effective quark propagator. The dispersion relation of the gluons in the plasma follows from the pole of this effective propagator. Due to the complicated momentum dependence of the HTL gluon self energy the dispersion relation can be obtained only numerically. Besides the branch belonging to transverse gluons, a plasmon branch shows up, describing the propagation of a longitudinal gluon in the plasma. The plasma frequency, defined as the zero momentum limit of the dispersion relation, is given by $m_g^* = gT/\sqrt{3}$. Instead of using the complicated dispersion relation, the transverse gluons have been regarded

as quasiparticles with the plasma frequency as effective mass. The plasmons, on the other hand, have been neglected as they are suppressed for typical momenta of the order of the temperature [10]. Although the effective masses have been calculated under the assumption of weak coupling ($g < 1$), the resulting equation of state is in good agreement with the one found on the lattice, even close to the phase transition where the coupling constant is not small.

Here we want to apply the idea of an ideal gas of quasiparticles with effective masses to the case of cold (strange) quark matter. Instead of gluons at finite temperature we consider up, down, and strange quarks at zero temperature but finite chemical potential μ . The effective quark mass is obtained from the zero momentum limit of the quark dispersion relation, which follows from the hard dense loop (HDL) approximation [11] of the quark self energy at finite chemical potential, where the internal momenta of the one-loop self energy diagram are assumed to be hard, i.e. of the order μ or larger. This calculation will be presented in the next section, where we focus in particular on the role of the current quark masses. In the third section we will discuss the statistical mechanics of a system of quasi-particles with density dependent masses. The effects of these masses on the equation of state of quark matter and a comparison with previous results without medium effects will be the topic of section 4. Finally we will apply our results to quark stars in section 5.

2 Medium-dependent effective quark masses

The aim of the present section is the derivation of an effective quark mass in an extended, dense system of quarks at zero temperature. This effective mass is defined as the zero momentum limit of the dispersion relation following from the quark self energy, which can be written in the most general form as [12]

$$\Sigma = -a P_\mu \gamma^\mu - b \gamma_0 - c, \quad (1)$$

where

$$\begin{aligned} a &= \frac{1}{4p^2} [tr(P_\mu \gamma^\mu \Sigma) - p_0 tr(\gamma_0 \Sigma)], \\ b &= \frac{1}{4p^2} [P^2 tr(\gamma_0 \Sigma) - p_0 tr(P_\mu \gamma^\mu \Sigma)], \\ c &= -\frac{1}{4} tr \Sigma, \end{aligned} \quad (2)$$

and $P^2 = p_0^2 - p^2$.

The dispersion relation is determined from the poles

of the effective propagator

$$S^*(P) = \frac{1}{P_\mu \gamma^\mu - m - \Sigma}, \quad (3)$$

leading to

$$[p_0(1+a) + b]^2 = p^2(1+a)^2 + (m-c)^2, \quad (4)$$

where m is the current quark mass. The traces over Σ are calculated in the hard dense loop (HDL) approximation [11], where only hard momenta of the order of the quark chemical potential μ for the internal quark and gluon lines in the one-loop quark self energy of fig. 1 are used. As in the high temperature case [9] the vacuum contribution and the contribution from integration over soft momenta are of higher order. This approximation is equivalent to the leading term in the high density approximation analogously to the finite temperature case [8]. In this way we arrive at a gauge invariant expression for the self energy, complete to leading order in the coupling constant g . Of course, due to the perturbative

nature of this approximation, the resulting expressions are only valid for small coupling constants, corresponding to a large chemical potential μ , i.e. high density, according to asymptotic freedom. However, as shown in the case of a hot gluon gas, the equation of state using effective masses calculated perturbatively agrees very well with lattice results even at temperatures where the coupling constant is not small [7]. (As a matter of fact, there are indications that for large coupling constants the lowest order results are better than higher order corrections [13].)

In the case of a vanishing current quark mass, as it can be assumed for up and down quarks, the HDL quark self energy is given by [8, 14]

$$\begin{aligned} \text{tr}\Sigma &= 0, \\ \text{tr}(P_\mu \gamma^\mu \Sigma) &= 4m_q^{*2}, \\ \text{tr}(\gamma_0 \Sigma) &= 2m_q^{*2} \frac{1}{p} \ln \frac{p_0 + p}{p_0 - p}, \end{aligned} \quad (5)$$

where

$$m_q^{*2} = \frac{g^2 \mu^2}{6\pi^2}. \quad (6)$$

(In the finite temperature case the only change is the replacement of m_q^{*2} by $g^2(T^2 + \mu^2/\pi^2)/6$ [14].)

Resumming this quark self energy leads to an effective propagator (3) which is part of a systematic expansion (Braaten-Pisarski method) at finite temperature and chemical potential [9, 14]. (For a review of this method as well as its applications to the quark-gluon plasma see [15].)

For non-vanishing current quark masses, e.g. for strange quarks with $m_s \simeq 150$ MeV, we have two possibilities for incorporating the current mass into the effective one. First the current mass could be hard, i.e. of the order of μ . In this case we have a second hard scale besides the chemical potential. Then the HDL approximation for the quark self energy may not be sufficient, as terms of the order of gm_s in the quark self energy from the integration over soft momenta or from the vacuum contribution may appear. Also gauge invariance of the self energy is not longer guaranteed. On the other hand, if we assume the current mass to be soft, $m_s \sim g\mu$, it can be neglected in the calculation of the quark self energy, where only hard momenta and energies are needed. The soft mass assumption might be justified, even for small g since $\mu \simeq 350$ MeV holds, as we will see below. Hence we end up with the same quark self energy as in the massless case (5) and (6).

Inserting (5), (6), and (2) into (4) in the massless case leads to the dispersion relation shown in fig. 2. The upper branch ω_+ corresponds to a collective quark mode, whereas the lower branch ω_- describes the propagation of a so-called plasmino [16], which is absent in the vacuum analogously to the plasmon solution for gluons. The effective mass at $p = 0$ is given by m_q^* . In the case of a current mass of the order of $m_s \sim g\mu$ we must

not neglect this mass in (3) and (4) since m_s and Σ are of the same order. In the zero momentum limit we find from (4) the effective strange quark mass [12]

$$m_s^* = \frac{m_s}{2} + \sqrt{\frac{m_s^2}{4} + \frac{g^2 \mu^2}{6\pi^2}}. \quad (7)$$

For evaluating the equation of state for an ideal gas of collective quarks, we neglect the plasmino as it is suppressed, i.e., the spectral strength function (residue of the pole of the effective propagator) vanishes exponentially for hard momenta of the order of μ [17] which dominate the equation of state. Furthermore we replace the dispersion relation of the collective quark branch by

$$\omega_{q,s}^*(k) = \sqrt{k^2 + m_{q,s}^{*2}}. \quad (8)$$

The simple dispersion relation ω_q^* is shown in fig. 2. It deviates from the exact one ω_+ by less than 11% over the entire momentum range. The deviation of ω_s^* from the exact solution can be shown to be even smaller.

Since there are no lattice calculation at finite density to which we could compare our results, the simple dispersion relation (8) is sufficient for our purpose, describing the role of medium effects in SQM at least qualitatively. Therefore we neglect further improvements of the quasiparticle treatment introduced in the case of a hot gluon gas [7] as the use of an effective number of degrees of freedom, of a temperature dependent coupling constant, and of an additional constant in front of the effective mass in the dispersion relation for fitting the high momentum part of the exact dispersion relation.

Finally we will discuss the value of g to be taken here. In principle it should be determined as a μ -dependent running coupling constant from the renormalisation group equation at finite density. However, so far there are no clear results at finite temperature or density [18]. Assuming the zero density result

$$\alpha_s(Q) = \frac{g^2}{4\pi} = \frac{12\pi}{(33 - 2n_f) \ln(Q^2/\Lambda^2)} \quad (9)$$

and replacing Q by the average three momentum in quark matter $\langle q \rangle = 3\mu/4$ we find for three quark flavors ($n_f = 3$), $\mu = 350$ MeV (see below), and the QCD scale parameter $\Lambda = 200$ MeV a value of $g = 5.7$. In calculations of SQM including lowest order α_s -corrections typical values of $g = 5.3$ ($\alpha_s = 2.2$) have been used [4], derived from comparing the MIT bag model with hadronic properties. In the approach proposed here one expects that g ought to be small as the effective mass is calculated perturbatively. However, in the case of a gluon gas the use of the lowest order effective mass provided an excellent agreement with lattice results even at the phase transition, where $g = 7.7$ was used [7]. As we will see below, an upper limit for the coupling constant in our approach is given by $g = 7.7$. As medium

effects in quark matter cannot be related directly to α_s -corrections fitted to hadronic properties and the zero density result of the running coupling constant cannot

be trusted, we take g as a free parameter ranging from 0 to 5.

3 Statistical mechanics of systems with medium dependent effective quark masses

The underlying philosophy to do statistical mechanics with medium dependent effective quark masses is to treat an ensemble of quasiparticles as a free, (in this case) degenerate Fermi gas of particles with a total energy

$$\hat{H}_{eff} = \sum_{i=1}^d \sum_{\mathbf{k}} \sqrt{\mathbf{k}^2 + m_i^{*2}(\mu)} \hat{a}_{\mathbf{k},i}^\dagger \hat{a}_{\mathbf{k},i} + E^*(\mu). \quad (10)$$

Here d denotes the degree of degeneracy (e.g. $d = 6n_f$ for n_f flavors). Due to the μ -dependence of $m^*(\mu)$ the function $E^*(\mu) = B^*(\mu)V$ defines a necessary energy counterterm in order to maintain thermodynamic self-consistency [19]. From this Hamiltonian the particle density ρ , energy density ϵ and pressure p at temperature $T = 0$ of a Fermi gas of free quasiparticles are given by

$$\begin{aligned} \rho(\mu) &= \frac{d}{2\pi^2} \int_{k=0}^{k_F} dk k^2 \\ &= \frac{d}{6\pi^2} k_F^3, \end{aligned} \quad (11)$$

$$\begin{aligned} \epsilon(\mu) - B^*(\mu) &= \frac{d}{2\pi^2} \int_{k=0}^{k_F} dk k^2 \omega^*(k) \\ &= \frac{d}{16\pi^2} \left[\mu k_F (2\mu^2 - m^{*2}) - m^{*4} \ln \left(\frac{k_F + \mu}{m^*} \right) \right], \end{aligned} \quad (12)$$

$$\begin{aligned} p(\mu) + B^*(\mu) &= \frac{d}{2\pi^2} \int_{k=0}^{k_F} dk k^2 (\mu - \omega^*(k)) \\ &= \frac{d}{24\pi^2} \left[\mu k_F \left(\mu^2 - \frac{5}{2} m^{*2} \right) + \frac{3}{2} m^{*4} \ln \left(\frac{k_F + \mu}{m^*} \right) \right]. \end{aligned} \quad (13)$$

Up to the additional function B^* , which can be regarded as a μ -dependent bag constant, these are the ideal Fermi gas formulas at temperature $T = 0$ for quasiparticles of mass m^* and chemical potential μ . The Fermi momentum is $k_F = (\mu^2 - m^{*2})^{1/2}$. The function $B^*(\mu)$ is necessary to maintain thermodynamic self-consistency. For example the fundamental thermodynamic relation between energy density and pressure

$$\epsilon(\mu) = \mu \frac{dp(\mu)}{d\mu} - p(\mu) \quad (14)$$

has to be fulfilled. So we should require additionally [19]

$$\left(\frac{\partial p}{\partial m^*} \right)_\mu = 0. \quad (15)$$

By using (13) we then find

$$\frac{dB^*}{dm^*} = -\frac{d}{4\pi^2} \left[m^* \mu k_F - m^{*3} \ln \left(\frac{k_F + \mu}{m^*} \right) \right]. \quad (16)$$

To carry out the integration of (16) we have to use the explicit functional dependence $m^*(\mu)$ respectively $\mu = \mu(m^*)$ in the two cases of light- and strange quarks.

3.1 EOS of massless quarks

In the case of $m^*(\mu) = m_q^*(\mu)$ (6) we integrate (16) and obtain

$$B_q^*(\mu_q) = -\frac{d_q}{16\pi^2} \left[\alpha^2 \beta - \alpha^4 \ln \left(\frac{\beta + 1}{\alpha} \right) \right] \mu_q^4, \quad (17)$$

choosing $B_q^*(0) = 0$. (Later we will add the phenomenological bag constant B_0 .) The g dependent functions $\alpha(g)$ and $\beta(g)$ are given by

$$\alpha(g) = \frac{g}{\sqrt{6\pi}}, \quad (18)$$

$$\beta(g) = \sqrt{1 - \alpha^2(g)}. \quad (19)$$

After inserting (17) into (11)-(13) and some manipulations using $m^* = \alpha(g)\mu$, the equations of state then assume the simple form

$$\rho_q(\mu_q) = \frac{d_q}{6\pi^2} \beta^3 \mu_q^3, \quad (20)$$

$$\epsilon_q(\mu_q) = \frac{d_q}{8\pi^2} \beta^3 \mu_q^4, \quad (21)$$

$$p_q(\mu_q) = \frac{d_q}{24\pi^2} \beta^3 \mu_q^4, \quad (22)$$

3.2 EOS of massive quarks

Taking $m^*(\mu)$ in (16) to be the effective strange quark mass $m_s^*(\mu_s)$ given by (7), the integration leads to ($d_s = 6$)

$$\begin{aligned} B_s^*(\mu_s) = & -\frac{d_s}{16\pi^2} \left[\frac{\sqrt{(m_s^* - m_s)(\beta^2 m_s^* - m_s)}}{24\alpha^2 \beta^4} \right. \\ & \times \sum_{n=0}^3 a_n m_s^{3-n} m_s^{*n} \\ & + \frac{5\alpha^4 - 12\alpha^2 + 8}{16\beta^5} m_s^4 \\ & \times \ln \left(\frac{\beta \sqrt{m_s^* - m_s} + \sqrt{\beta^2 m_s^* - m_s}}{\alpha^2 m_s} \right) \\ & \left. - m_s^{*4} \ln \left(\frac{k_F + \mu}{m_s^*} \right) \right], \end{aligned} \quad (23)$$

which obviously fulfill $\rho_q = \partial p_q / \partial \mu_q$ and $\epsilon_q + p_q = \mu_q \rho_q$. They are the well known equations for massless fermions up to the factor β^3 ($0 < \beta(g) < 1$). Note that ρ_q , ϵ_q and p_q decreases with increasing coupling constant g . On the other hand the implicit form $p_q = p_q(\epsilon_q)$ of the EOS $p_q = \epsilon_q/3$ does not reveal any g dependence, while the EOS in the form $p_q = p_q(\rho_q) \propto \rho_q^{4/3}/\beta(g)$ slightly depends on g . The EOS in the form $p_q = p_q(\epsilon_q)$ (with respect to modifications due to massive s -quarks and electrons) will be important for the investigations of SQM-stars in section 5. From (19) we observe that g must not exceed the value $g = \sqrt{6\pi} \simeq 7.7$.

To get an idea of the order of magnitude of B_q^* (which is negative), fig. 3 shows $|B_q^*|^{1/4}$ versus μ_q for various values of g . If $\mu \approx 350$ MeV holds (see below) $|B_q^*|^{1/4}$ is of the order of 100 MeV. Using (17) and (21) this corresponds to $B_q^*/\epsilon_q < 10\%$ for $g < 5$.

where

$$\begin{aligned} a_0(g) &= -15\alpha^4 + 26\alpha^2 - 8, \\ a_1(g) &= -10\alpha^4 + 18\alpha^2 - 8, \\ a_2(g) &= -8\alpha^4 + 16\alpha^2 - 8, \\ a_3(g) &= 24\alpha^4 - 48\alpha^2 + 24. \end{aligned}$$

From requiring a positive Fermi momentum, $k_F = (\mu^2 - m^{*2})^{1/2} \geq 0$, we find $\mu_s \geq m_s^*$, from which $\mu_s \geq m_s \beta^{-2}$ follows. This constraint is equivalent to $\beta^2 m_s^* \geq m_s$ needed in (23). At $\mu_s = m_s \beta^{-2}$ the integration constant is chosen in such a way that B_s^* vanishes. Utilizing (23) in connection with (11)-(13) (using $m^* = m_s^*$) leads to the equations of state $\rho_s(\mu_s)$, $\epsilon_s(\mu_s)$ and $p_s(\mu_s)$ for strange quarks.

4 Medium-effects in strange quark matter

In the following sections we investigate in detail within our model the equation of state and the properties of SQM.

4.1 Strange quark matter consisting of massless quarks

It is illustrative to consider first the case of SQM consisting of massless quarks ($m_{u,d,s} = 0$). In section 3.1 we derived the equations of state (20)-(22) for light quarks with the effective quark mass (6). We saw that ρ_q , ϵ_q and p_q decreases with increasing coupling constant g . Now we want to investigate how this affects the energy

per baryon (E/A) of massless SQM

$$\left(\frac{E}{A} \right) (\mu_q) = \frac{\epsilon_q(\mu_q) + B_0}{\rho_B(\mu_q)} \quad (24)$$

$$= 3 \left(\frac{3}{4} \mu_q + \frac{6\pi^2}{d_q \beta^3 \mu_q^3} B_0 \right).$$

Here B_0 denotes the usual bag-constant which corresponds to the energy difference between the "false", perturbative vacuum inside the bag and the "true" vacuum at the outside, and which, in return, confines the quarks. The degree of degeneracy is given by $d_q = 6n_f = 18$ and $\rho_B(\mu_q) = \rho_q(\mu_q)/3$ is the baryon density. The minimum energy with respect to μ_q is found to be

$$\left(\frac{E}{A} \right)_{min} = 3\mu_{min}, \quad (25)$$

where

$$\mu_{min} = \left(\frac{24\pi^2}{d_q \beta^3} B_0 \right)^{1/4} \quad (26)$$

is the chemical potential which minimizes E/A . The optimum chemical potential μ_{min} ranges from 276 MeV for $g = 0$ and $B_0^{1/4} = 145$ MeV to 374 MeV for $g = 5$ and $B_0^{1/4} = 160$ MeV. Minimizing the energy is here equivalent to the equilibrium condition $p = p_q - B_0 = 0$ between the Fermi pressure inside the bag and the pressure of the nonperturbative QCD vacuum outside the bag given by B_0 . This leads to an electrically neutral

4.2 Strange quark matter

Now we use the equations of state derived in section 3.1 and 3.2 to calculate the energy per baryon of strange quark matter ($d_q = 12$, $d_s = 6$) assuming finite s -quark masses:

$$\left(\frac{E}{A} \right) (\mu_q, \mu_s) = \frac{\epsilon_q(\mu_q) + \epsilon_s(\mu_s) + B_0}{\rho_B(\mu_q, \mu_s)}. \quad (27)$$

The baryon density is now given by

$$\rho_B(\mu_q, \mu_s) = \frac{\rho_q(\mu_q) + \rho_s(\mu_s)}{3}. \quad (28)$$

As usual we define the strangeness fraction f_s to be

$$f_s(\mu_q, \mu_s) = \frac{\rho_s(\mu_s)}{\rho_B(\mu_q, \mu_s)}, \quad (29)$$

yielding the number of s -quarks per baryon. Fig. 4 shows the minimum energy per baryon E/A (i.e. $p = p_u + p_d + p_s - B_0 = 0$) for a given strangeness content f_s and various values of g . The bag constant is assumed to be $B_0^{1/4} = 145$ MeV. The current s -quark mass is $m_s = 150$ MeV. One sees that E/A increases with increasing g . At $g = 0$ the *absolute* minimum is achieved for a strangeness fraction of $f_s \approx 0.7$. The absolute minimum corresponds to a system where all flavors assume the same Fermi energy (i.e. $\mu_q = \mu_s$).

system with an equal number of u , d and s -quarks. Eq. (25) shows that in the case of massless quark matter the medium effects, we take into account, lead to an increase of the energy per baryon by the factor

$$\mu_{min} \propto \beta^{-3/4} = \left(1 - \frac{g^2}{6\pi^2} \right)^{-3/8}.$$

For instance assuming $B_0^{1/4} = 145$ MeV ($B_0 = 58$ MeV/fm³) this increases $(E/A)_{min}$ from about 829 MeV ($g = 0$) to 881 MeV ($g = 3$, $\alpha_s = 0.72$). That is to say, owing to medium effects the binding energy per baryon of the system with respect to ^{56}Fe is reduced from about 100 MeV to 50 MeV. The more massive constituents raise the overall energy scale.

At $p = 0$ ($\mu = \mu_{min}$) we find using (20)-(22) that

$$\rho_q(\mu_{min}) \propto \beta^3 \mu_{min}^3 \propto \beta^{3/4}$$

decreases while

$$\epsilon_q(\mu_{min}) \propto \beta^3 \mu_{min}^4 \propto \beta^3 \beta^{-3} = \text{const.}$$

is independent of g .

In the following we focus on the properties of SQM in its *absolute* minimum. Fig. 5 shows this minimal energy versus g . Assuming $m_s = 150$ MeV and $B_0^{1/4} = 145$ MeV the minimal energy increases from 874 MeV ($g = 0$) to 943 MeV ($g = 3$). Therefore, for the chosen parameters SQM is not bound with respect to ^{56}Fe . As already seen in the massless case ($m_s = 0$) the baryon density at $p = 0$ decreases with increasing g (fig. 6) while the energy density ϵ stays approximately constant (fig. 7). The strangeness fraction f_s is found to decrease with increasing g (fig. 8) which is in contrast to the massless calculation where $f_s = 1$ due to flavor symmetry. Furthermore, this result is also in contrast to the effect of α_s -corrections on f_s which will be discussed in the next section. The decreasing strangeness fraction does not seem to be an artifact of the special form of the effective mass m_s^* given by (7). For instance the same behaviour holds for a "hypothetical" mass of the form

$$m_s^* = m_s + \frac{g\mu}{\sqrt{6}\pi}.$$

So, in this approach medium effects lead to less dense and less bound strange quark matter with a lowered number of strange quarks per baryon!

4.3 Medium effects versus bag constant

As shown above, describing SQM with medium effects using effective quark masses leads to the mass formula (27) depending on the additional parameter g . From a practical point of view, introducing the new parameter g gives rise to the question whether the effect of changing g , say the effect of the medium, can already be described by a variation of the original parameters that enter the model, namely B_0 and m_s . The bag constant might be suitable for this purpose because of the uncertainty of its value and its strong influence on the behavior of SQM (figs. 5-8). Therefore we will briefly discuss whether a change in the bag constant ($B_0 \rightarrow B_0 + \Delta B_0$) at $g = 0$ is able to approximately recover changes of the properties (E/A , ρ , ϵ) due to medium effects parameterized by g . As a criterion to compare the "medium" approach (where B_0 is unchanged and $g > 0$) with the "bag" approach (changing B_0 at $g = 0$) we require that both should lead to the same energy per baryon at its absolute minimum, i.e.

$$(E/A)_{med}(B_0, g > 0) \stackrel{!}{=} (E/A)_{bag}(B_0 + \Delta B_0, g = 0). \quad (30)$$

Here "med" and "bag" stands for the "medium" and the "bag" approach, respectively. Under this condition we calculate ρ_{med} , ϵ_{med} for different values of g and compare it to the associated ρ_{bag} , ϵ_{bag} at a ΔB_0 so that (30) holds. Fig. 9 and fig. 10 show the minimal energy for various f_s (see figure caption) versus ρ and ϵ . One sees that both approaches lead to a different physical behavior. Whereas ρ_{med} , ϵ_{med} decreases (or stays constant) with increasing g , one sees that ρ_{bag} , ϵ_{bag} strongly increases. The same different behavior appears for f_s . From this we can conclude that it is not possible to describe the influence of medium effects just by altering the bag constant. For completeness, we should mention that similar arguments hold for the variation of m_s instead of B_0 .

Furthermore, the EOS of SQM has been computed including α_s -corrections, i.e. one-gluon exchange [3, 21]. Increasing α_s there leads to an increase of E/A as in our approach. However, at the same time f_s increases in contrast to our case. Hence, we conclude that the influence of the effective quark masses on SQM cannot be simulated by taking into account α_s -corrections or changing B_0 or m_s .

5 Application to strange quark matter stars

Originally SQM in bulk was thought to exist only in the interior of neutron stars where the pressure is high enough that the neutron matter melts into its quark substructure [22]. At least in the cores of neutron stars, where the density rises up to the order of 10 times normal nuclear density, it is not very likely that matter consists of individual hadrons. It thus still might be that SQM exists in neutron stars. For an overview of the structure of neutron stars, see Ref. [23].

In this section we investigate neutron stars under the assumption that they *entirely* consist of cold electrically charge neutral SQM (SQM stars) in equilibrium with respect to the weak interaction. This requires the inclusion of electrons into the model. We focus on the change of the properties of SQM-stars ($p(\epsilon)$, M , R , E/A) due to medium effects in the spirit of the previous sections. Here M and R denotes the mass and radius of the star.

The thermodynamic of electrons at $T = 0$ is given by (11)-(13) taking $d = 2$ and $m^* = 0$

$$\rho_e = \frac{\mu_e^3}{3\pi^2}, \quad (31)$$

$$\epsilon_e = \frac{\mu_e^4}{4\pi^2}, \quad (32)$$

$$p_e = \frac{\mu_e^4}{12\pi^2}. \quad (33)$$

Now, we are dealing with four chemical potentials (μ_u , μ_d , μ_s , μ_e) which are related by the chemical equilib-

rium between the quark flavors and the leptons. The basic weak reactions are given by

$$d \longrightarrow u + e^- + \bar{\nu}_e \quad (34)$$

$$s \longrightarrow u + e^- + \bar{\nu}_e. \quad (35)$$

The equilibration of flavors is provided by [24]

$$s + u \longrightarrow d + u. \quad (36)$$

Hence, the four chemical potentials are reduced to two independent ones

$$\mu \equiv \mu_s = \mu_d \quad \text{and} \quad \mu_u = \mu - \mu_e. \quad (37)$$

Finally, the condition of electrically charge neutrality

$$2/3\rho_u - 1/3(\rho_d + \rho_s) - \rho_e = 0 \quad (38)$$

just leaves one independent chemical potential, say μ . Therefore, the EOS can be written as a function of μ only:

$$\rho_B = (\rho_u + \rho_d + \rho_s)/3, \quad (39)$$

$$\epsilon = \epsilon_u + \epsilon_d + \epsilon_s + \epsilon_e + B_0, \quad (40)$$

$$p = p_u + p_d + p_s + p_e - B_0, \quad (41)$$

where $\rho_{u,d}$, $\epsilon_{u,d}$ and $p_{u,d}$ are calculated from (20)-(22) with $d_q = 6$ and individual chemical potentials μ_u , μ_d .

One of the aim of this section is to determine whether there are significant changes in mass M and radius R of SQM stars due to medium effects. To do so, we assume a cold, static, spherical star. It is described by the solutions of the Oppenheimer-Volkoff-Tolman (OVT) equations of hydrostatic equilibrium. They follow from general relativity [25] which one has to apply due to highly concentrated matter and therefore curved space-time. The solutions of the OVT equations are the pressure $p(r)$ and energy density $\epsilon(r)$ at a given radius r inside the star. The radius R of the star is determined by the condition $p(R) = 0$ while the mass is given by

$$M = \int_{r=0}^R 4\pi r^2 \epsilon(r) dr. \quad (42)$$

The OVT equation can be solved specifying the central energy density $\epsilon_c = \epsilon(r=0)$ and the EOS in the form

6 Conclusions

In the present work we proposed an improved description of SQM by taking medium effects into account. We considered the quarks as quasiparticles, which acquire an effective mass generated by the interaction with other quarks of the dense system. The effective masses are derived from the zero momentum limit of the dispersion relations following from the quark self energy in the hard dense loop approximation. We regard the strong coupling constant g contained in the effective mass as a free parameter. Although the quark self energy has been calculated perturbatively we extrapolate our final results up to $g = 5$. The validity of this approach is supported by the success of an equivalent method applied to a hot gluon gas [7].

In the case of quark matter consisting of quarks with zero current mass the density (20), the energy density (21), and the pressure (22) are simply changed by a factor $\beta^3(g)$ (19), which decreases with increasing g starting from $\beta = 1$ at $g = 0$. As a consequence the energy per nucleon increases with increasing g . The EOS $\epsilon(p)$, on the other hand, turns out to be independent of g .

Acknowledgements

We would like to thank N. Glendenning, A. Peshier and F. Weber for helpful discussions. K.S. and M.H.T. are grateful to the University of Regensburg for their hospitality.

$p = p(\epsilon)$. The numerical calculation of $p = p(\epsilon)$ from (40), (41) shows that there is no noteworthy influence by a change of g . The EOS gets slightly softer for high values of the coupling constant (e.g. $g = 4$) but there is no remarkable impact on R and M (see fig. 11).

Note, that this result is only valid under the assumption of a pure SQM star. Although there is no change in mass and radius of the star, there is, nevertheless, an increase of the energy E/A at a given radius inside the star due to medium effects. This increases the possibility of a phase transition to hadronic matter and therefore the possibility of a nuclear crust on the surface of the SQM core. Figs. 12 and 13 show this behavior as a function of p and ϵ , respectively. In fig. 12 the pressure inside the star decreases from a given central pressure $p_c = p(\epsilon_c)$ in the center of the star to $p = 0$ on the surface where E/A is minimal. One finds that the energy E/A is increased in the entire star. Furthermore, fig. 14 shows the change of the strangeness fraction versus the baryon density for different values of g .

We conclude the discussion of strange stars with the result that mass and radius of pure SQM stars are not changed by medium effects while, considering hybrid stars, mass and radius of the star might be indirectly influenced by medium effects due to a change of the crust formation.

This picture holds qualitatively also for the case of SQM, where a non-vanishing current mass of the strange quark is considered. Even in the case of a small bag constant $B_0^{1/4} = 145$ MeV and a small strange mass of $m_s = 150$ MeV the energy per baryon E/A of SQM is larger than the one of ^{56}Fe already for $g = 3$. For higher values of m_s , $B_0^{1/4}$, or g even larger values for E/A are found. This suggests that SQM is less bound if medium effects are taken into account. However, it might still be metastable, i.e. stable against decays caused by the strong interaction, which then has interesting consequences for the formation of strangelets in ultrarelativistic heavy ion collisions [21].

Applying our results to strange quark stars, we observe that the radius-mass relation of these stars is hardly changed by the presence of an effective quark mass since the EOS depends only very weakly on it. However, owing to the increase of E/A with an increasing effective mass the phase transition to hadronic matter will take place at a smaller radius in the interior of the star leading to a thicker crust.

References

- [1] A.R. Bodmer, Phys. Rev. D 4 (1971) 1601; E. Witten, Phys. Rev. D 30 (1984) 272.
- [2] Strange Quark Matter in Physics and Astrophysics, edited by J. Madsen and P. Haensel, Nucl. Phys. B (Proc. Suppl.) 24B.
- [3] B.A. Freedman and L.D. McLerran, Phys. Rev. D 16 (1978) 1169.
- [4] S.A. Chin and A.K. Kerman, Phys. Rev. Lett. 43 (1979) 1292.
- [5] E. Fahri and R.L. Jaffe, Phys. Rev. D 30 (1984) 2379.
- [6] V. Goloviznin and H. Satz, Z. Phys. C 57 (1993) 671; A. Peshier et al., Phys. Lett. B 337 (1994) 235.
- [7] A. Peshier, B. Kämpfer, O.P. Pavlenko and G. Soff, Phys. Rev. D 54 (1996) 2399.
- [8] V.V. Klimov, Sov. Phys. JETP 55 (1982) 199; H.A. Weldon, Phys. Rev. D 26 (1982) 1394 and 2789.
- [9] E. Braaten and R.D. Pisarski, Nucl. Phys. B 337 (1990) 310.
- [10] R.D. Pisarski, Physica A 158 (1989) 146.
- [11] C. Manuel, Phys. Rev. D 53 (1996) 5866.
- [12] R.D. Pisarski, Nucl. Phys. A 498 (1989) 423c; J.-P. Blaizot and J.-Y. Ollitrault, Phys. Rev. D 48 (1993) 1390.
- [13] A. Peshier, private communication.
- [14] H. Vija and M.H. Thoma, Phys. Lett. B 342 (1995) 212.
- [15] M.H. Thoma, Quark-Gluon Plasma 2, edited by R.C. Hwa, (World Scientific, Singapore, 1995) p.51.
- [16] E. Braaten, Astrophys. J. 392 (1992) 70; E. Braaten, R.D. Pisarski, and T.C. Yuan, Phys. Rev. Lett. 64 (1990) 2242.
- [17] H.A. Weldon, Phys. Rev. D 40 (1989) 2410; G. Baym, J.-P. Blaizot, and B. Svetitsky, Phys. Rev. D 46 (1992) 4043.
- [18] R. Baier, B. Pire, and D. Schiff, Z. Phys C 51 (1991) 581; M.A. van Eijck, C.R. Stephens, and C.G. van Weert, Mod. Phys. Lett A 8 (1994) 309.
- [19] M.I. Gorenstein and S.N. Yang, Phys. Rev. D 52 (1995) 5206.
- [20] J. D. Bjorken and L. D. McLerran, Phys. Rev. D 20 (1979) 2353.
- [21] C. Greiner, P. Koch and H. Stöcker, Phys. Rev. Lett. 58 (1987) 1825; C. Greiner, D.-H. Rischke, H. Stöcker and P. Koch, Phys. Rev. D 38 (1988) 2797; C. Greiner and H. Stöcker, Phys. Rev. D 44 (1991) 3517; for a recent review see C. Greiner and J. Schaffner, Quark-Gluon Plasma 2, edited by R.C. Hwa, (World Scientific, Singapore, 1995) p.635.
- [22] J. C. Collins and M. J. Perry, Phys. Rev. Lett. 34 (1975) 1353; G. Baym and S. Chin, Phys. Lett. 62 B (1976) 241; B. A. Freedman and L. D. McLerran, Phys. Rev. D 16 (1978) 1169.
- [23] S.L. Shapiro and S.A. Teukolsky, Black Holes, White Dwarfs, and Neutron Stars, (John Wiley & Sons, N.Y., 1983); F. Weber and N. K. Glendenning, astro-ph/9609074 (1996).
- [24] Ch. Kettner, F. Weber, M.K. Weigel and N.K. Glendenning, Phys. Rev. D 51 (1995) 1440.
- [25] J. R. Oppenheimer and G. M. Volkoff, Phys. Rev. 55 (1939) 374.




Figure 1: One-loop quark self energy




Figure 2: Quark dispersion relations




Figure 3: $B_q^*(\mu)$ (17) for various values of g ; $d_q = 18$ (assuming $m_s = 0$)




Figure 4: Energy per baryon versus the strangeness fraction at $B_0^{1/4} = 145$ MeV and $m_s = 150$ MeV




Figure 5: Energy per baryon versus the coupling constant at $m_s = 150$ MeV




Figure 6: Baryon density versus the coupling constant at $m_s = 150$ MeV, $\rho_0 = 0.17 fm^{-3}$




Figure 7: Energy density versus the coupling constant at $m_s = 150 \text{ MeV}$




Figure 8: Strangeness fraction versus the coupling constant at $m_s = 150 \text{ MeV}$




Figure 9: Comparison of ρ in the "medium" approach and the "bag" approach at $m_s = 150$ MeV and $B_0^{1/4} = 145$ MeV. The points denote the increase of f_s from $f_s = 0$ (top left) to $f_s = 1.5$ (top right) in steps of 0.1; $\rho_0 = 0.17 \text{ fm}^{-3}$




Figure 10: Comparison of ϵ in the "medium" approach and the "bag" approach at $m_s = 150$ MeV and $B_0^{1/4} = 145$ MeV. The points denote the increase of f_s from $f_s = 0$ (top left) to $f_s = 1.5$ (top right) in steps of 0.1




Figure 11: Mass (in units of solar mass) versus radius of a SQM star, $m_s = 150$ MeV, $B_0^{1/4} = 145$ MeV




Figure 12: E/A versus p in a SQM star, $m_s = 150$ MeV and $B_0^{1/4} = 145$ MeV




Figure 13: E/A versus ϵ in a SQM star, $m_s = 150$ MeV and $B_0^{1/4} = 145$ MeV




Figure 14: Strangeness fraction f_s versus ρ/ρ_0 in a SQM star, $m_s = 150$ MeV, $B_0^{1/4} = 145$ MeV and $\rho_0 = 0.17 \text{ fm}^{-3}$